


**AMENDMENTS TO THE CLAIMS**

Claims 1-21 (Canceled).

22. (Currently amended) ~~The 3,4-Alkylenedioxythiophenes of Claim 21,~~

~~wherein~~ A 3,4-Alkylenedioxythiophenes of the formula (I),




wherein

A is a C<sub>1</sub>-C<sub>5</sub>-alkylene radical which is substituted at any point by a linker L and optionally bears further substituents.


L is a methylene group.

p is 0 or an integer from 1 to 6.

M is an n-functional mesogenic group of the formula (II-a) or (II-b),




(II-a)




(II-b)

wherein


$X^1, X^2, X^3$ ,  $X^1, X^2$  and  $X^3$  are substituted or unsubstituted structures selected independently from the group consisting of





and

$Z^1, Z^2$   $Z^1$  and  $Z^2$  are structures selected independently from the group consisting of



wherein

$R^x$  and  $R^y$  are each, independently of one another, H, substituted or unsubstituted  $C_1$ - $C_{22}$ -alkyl,  $C_1$ - $C_{22}$ -haloalkyl,  $C_1$ - $C_{22}$ -alkenyl,  $C_1$ - $C_{22}$ -alkoxy,  $C_1$ - $C_{22}$ -thioalkyl,  $C_1$ - $C_{22}$ -iminoalkyl,  $C_1$ - $C_{22}$ -alkoxycarbonyl,  $C_1$ - $C_{22}$ -alkoxycarbonyloxy, a radical of an

aliphatic C<sub>1</sub>-C<sub>22</sub>-alkanecarboxylic acid or of acrylic acid, halogen, pseudohalogen, NO<sub>2</sub>, a carboxyl group or a hydroxy group,

h is an integer from 1 to 10,

w is an integer from 1 to 5,

~~x, y, z~~ x, y and z are each, independently of one another, 0 or 1, and

n is an integer from 1 to 8 and 1 or 2, where when n is 1, the group of the formula (II-a) or (II-b) bears a terminal ~~group F~~ group G at the linkage points denoted by \*,


~~wherein~~

~~F~~

wherein

G is substituted or unsubstituted C<sub>1</sub>-C<sub>22</sub>-alkyl, C<sub>1</sub>-C<sub>22</sub>-haloalkyl, C<sub>1</sub>-C<sub>22</sub>-alkenyl, C<sub>1</sub>-C<sub>22</sub>-alkoxy, C<sub>1</sub>-C<sub>22</sub>-thioalkyl, C<sub>1</sub>-C<sub>22</sub>-iminoalkyl, C<sub>1</sub>-C<sub>22</sub>-alkoxycarbonyl, C<sub>1</sub>-C<sub>22</sub>-alkoxycarbonyloxy, a radical of an aliphatic C<sub>1</sub>-C<sub>22</sub>-alkanecarboxylic acid or of acrylic acid, halogen, pseudohalogen, a nitro (NO<sub>2</sub>) group, a carboxyl group, a sulphonic acid group or sulphonate group or a hydroxy group,

B is a bridging group of the formula (B)



(B)

wherein

q is 0 or 1,

r and s are each 0 or 1, with the proviso that when r is 1, s is 0 and vice versa or both are optionally 0,


t is 0 or 1,

Sp is a spacer selected from the group consisting of substituted and unsubstituted linear or cyclic  $C_1$ - $C_{20}$ -alkylene groups,  $C_5$ - $C_{20}$ -arylene groups,  $C_2$ - $C_{20}$ -heteroarylene groups in which from one to three heteroatoms selected from the group consisting of N, O and S can additionally be present in the heteroaromatic ring or ring system,  $C_6$ - $C_{20}$ -aralkylene groups,  $C_2$ - $C_{200}$ -oligoether and -polyether groups,

m is 0 or 1,


Q is O, S or NH

with the proviso that said polythiophenes is not



23. (Currently amended) The 3,4-Alkylenedioxythiophenes of ~~Claim 21~~ claim 22,  
wherein

M is an n-functional mesogenic group selected from the group consisting of the formulae (II-c-1) to (II-c-6),



wherein

n is at most 4, 6 or 8,


and wherein when n is an integer below 4, 6 or 8, M is selected from the group consisting of the formulae (II-c-1) to (II-c-6) bearing a terminal group F on the remaining 4 - n, 6 - n or 8 - n linkage points denoted by \*,

wherein


F is H, substituted or unsubstituted C<sub>1</sub>-C<sub>22</sub>-alkyl, C<sub>1</sub>-C<sub>22</sub>-haloalkyl, C<sub>1</sub>-C<sub>22</sub>-alkenyl, C<sub>1</sub>-C<sub>22</sub>-alkoxy, C<sub>1</sub>-C<sub>22</sub>-thioalkyl, C<sub>1</sub>-C<sub>22</sub>-iminoalkyl, C<sub>1</sub>-C<sub>22</sub>-

alkoxycarbonyl, C<sub>1</sub>-C<sub>22</sub>-alkoxycarbonyloxy, a radical of an aliphatic C<sub>1</sub>-C<sub>22</sub>-alkanecarboxylic acid or of acrylic acid, halogen, pseudohalogen, a nitro (NO<sub>2</sub>) group, a carboxyl group, a sulphonic acid group or sulphonate group or a hydroxy group.

24. (Currently amended) The 3,4-Alkylenedioxythiophene of ~~Claim 21~~ claim 22, having the structure of the formulae (I-a) and/or (I-b),




(I-a)




(I-b)

25. (Currently amended) A mixture comprising at least one compound of formulae (I-a) and/or (I-b) according to ~~Claim 21~~:



(I-a)




(I-b)

wherein

M is an n-functional mesogenic group,

B is a bridging group of the formula (B)





(B)

wherein

$q$  is 0 or 1,

$r$  and  $s$  are each 0 or 1, with the proviso that when  $r$  is 1,  $s$  is 0 and vice versa or both are optionally 0,


$t$  is 0 or 1,

$\text{Sp}$  is a spacer selected from the group consisting of substituted and unsubstituted linear or cyclic  $\text{C}_1\text{-C}_{20}$ -alkylene groups,  $\text{C}_5\text{-C}_{20}$ -arylene groups,  $\text{C}_2\text{-C}_{20}$ -heteroarylene groups in which from one to three heteroatoms selected from the group consisting of N, O and S can additionally be present in the heteroaromatic ring or ring system,  $\text{C}_6\text{-C}_{20}$ -aralkylene groups,  $\text{C}_2\text{-C}_{200}$ -oligoether and -polyether groups,


$m$  is 0 or 1,

$\text{Q}$  is O, S or NH

with the proviso that said polythiophenes is not



26. (Previously presented) A 3,4-Alkylenedioxythiophene of the formula (I),



(I)

wherein

A is a C<sub>1</sub>-C<sub>5</sub>-alkylene radical which is substituted at any point by a linker L and optionally bears further substituents,


L is a methylene group,

p is 0 or an integer from 1 to 6,

M is an n-functional steroid radical or a derivative of a steroid radical,

n is an integer from 1 to 8 and

B is a bridging group of the formula (B)



(B)

wherein

q is 0 or 1,

r, s are each 0 or 1, with the proviso that when r is 1, s is 0 and vice versa or both are optionally 0,


t is 0 or 1,

Sp is a spacer selected from the group consisting of substituted and unsubstituted linear or cyclic C<sub>1</sub>-C<sub>20</sub>-alkylene groups, C<sub>5</sub>-C<sub>20</sub>-arylene groups, C<sub>2</sub>-C<sub>20</sub>-hetero-arylene groups in which from one to three heteroatoms selected from the group consisting of N, O and S can additionally be present in the heteroaromatic ring or ring system, C<sub>6</sub>-C<sub>20</sub>-aralkylene groups, C<sub>2</sub>-C<sub>200</sub>-oligoether and -polyether groups,

m is 0 or 1,


Q is O, S or NH.

27. (Currently amended) A 3,4-Alkylenedioxythiophene of the formula (I),



wherein


- A is a C<sub>1</sub>-C<sub>5</sub>-alkylene radical which is substituted at any point by a linker L and optionally bears further substituents,
- L is a methylene group,
- p is 0 or an integer from 1 to 6,
- M is an n-functional cholesteryl radical or a derivative of the cholesteryl radical of the formula (III-a),



(III-a)

wherein R is H, substituted or unsubstituted C<sub>1</sub>-C<sub>22</sub>-alkyl, C<sub>1</sub>-C<sub>22</sub>-haloalkyl, C<sub>1</sub>-C<sub>22</sub>-alkenyl, C<sub>1</sub>-C<sub>22</sub>-alkoxy, C<sub>1</sub>-C<sub>22</sub>-thioalkyl, C<sub>1</sub>-C<sub>22</sub>-iminoalkyl, C<sub>1</sub>-C<sub>22</sub>-alkoxycarbonyl, C<sub>1</sub>-C<sub>22</sub>-alkoxycarbonyloxy, a radical of an aliphatic C<sub>1</sub>-C<sub>22</sub>-alkanecarboxylic acid or of acrylic acid, halogen, pseudohalogen, a nitro (NO<sub>2</sub>) group, a carboxyl group, a sulphonic acid group or sulphonate group or a hydroxy ~~group-group~~.


- n is 1 and
- B is a bridging group of the formula (B)



wherein

- q is 0 or 1,  
 r, s are each 0 or 1, with the proviso that when r is 1, s is 0 and vice versa or both are optionally 0,  
 t is 0 or 1,  
 Sp is a spacer selected from the group consisting of substituted and unsubstituted linear or cyclic C<sub>1</sub>-C<sub>20</sub>-alkylene groups, C<sub>5</sub>-C<sub>20</sub>-arylene groups, C<sub>2</sub>-C<sub>20</sub>-hetero-arylene groups in which from one to three heteroatoms selected from the group consisting of N, O and S can additionally be present in the heteroaromatic ring or ring system, C<sub>6</sub>-C<sub>20</sub>-aralkylene groups, C<sub>2</sub>-C<sub>200</sub>-oligoether and -polyether groups,  
 m is 0 or 1,  
 Q is O, S or NH.

28. (currently amended) A process for ~~preparing~~ preparing a polythiophene comprising polymerizing the 3,4-alkylenedioxythiophene as claimed in claim 22 of the formula (I),




wherein

- A is a C<sub>1</sub>-C<sub>5</sub>-alkylene radical which is substituted at any point by a linker L and optionally bears further substituents,  
 L is a methylene group,  
 p is 0 or an integer from 1 to 6,

M — is an n-functional mesogenic group;

n — is an integer from 1 to 8 and

B — is a bridging group of the formula (B)



wherein

q — is 0 or 1;

r, s — are each 0 or 1, with the proviso that when r is 1, s is 0 and vice versa or both are optionally 0;


t — is 0 or 1;

Sp — is a spacer selected from the group consisting of substituted and unsubstituted linear or cyclic C<sub>4</sub>-C<sub>20</sub>-alkylene groups, C<sub>5</sub>-C<sub>20</sub>-arylene groups, C<sub>2</sub>-C<sub>20</sub>-hetero-arylene groups in which from one to three heteroatoms selected from the group consisting of N, O and S can additionally be present in the heteroaromatic ring or ring system, C<sub>6</sub>-C<sub>20</sub>-aralkylene groups, C<sub>2</sub>-C<sub>200</sub>-oligoether and polyether groups;

m — is 0 or 1;

Q — is O, S or NH.


with the proviso that said 3,4-Alkylenedioxythiophene is not




29. (Previously presented) The process of Claim 28 wherein a mixture of two or more compounds of Formula 1 are polymerized.

30. Cancelled

31. (Currently amended) The polythiophene according to claim 44, wherein the polythiophene  
~~Polythiophene according to Claim 30, characterized in that the comprise recurring units of~~  
 the formulae (IV-a) and/or (IV-b),



(IV-a)




(IV-b)

32. (Currently amended) ~~Polythiophene of Claim 30~~ The polythiophene of claim 44, wherein  
 they are cationically and electrically conductive and contain bound anions as counterions to  
 balance the positive charge.

33. (Previously presented) The polythiophene of Claim 32, wherein the counterions are  
 polyanions of polymeric carboxylic acids or polymeric sulphonic acids.

34. (Currently amended) The polythiophene according to ~~Claim 30~~ claim 44, wherein they are  
 uncharged and semiconducting.

35. (Currently amended) ~~The process for preparing polythiophenes of Claim 30, A process for~~  
preparing the polythiophene as claimed in claim 44, comprising oxidatively polymerizing  
electrochemically compounds of the formula (I),




(I)

36. (Currently amended) A process for preparing electrical or electronic components, light-emitting components, for antistatic coating, in optoelectronics or in solar energy technology comprising incorporating polythiophenes according to ~~Claim 30~~ claim 44.

37. (Currently amended) A process for preparing conductive layers comprising incorporating the polythiophenes of ~~Claim 30~~ claim 44.

38. (Cancelled)


39. (Currently amended) ~~Polythiophenes characterized in that they~~ A polythiophene which comprise recurring units of the formula (IV),



(IV)


produced according to the process of ~~Claim 38~~ claim 45

with the proviso that said polythiophenes is not



40. (Currently amended) A process for preparing electrical or electronic components, light-emitting components, for antistatic coating, in optoelectronics or in solar energy technology comprising incorporating the ~~polythiophenes~~ polythiophene of Claim 39.
41. (Currently amended) A process for preparing conductive layers comprising incorporating the ~~polythiophenes~~ polythiophene according to Claim 39.
42. (Previously presented) The process according to claim 37, which further comprises heating the layer at a temperature form 80<sup>0</sup>C to 300<sup>0</sup>C.
43. (Previously presented) The process according to claim 41, which further comprises heating the layer at a temperature form 80<sup>0</sup>C to 300<sup>0</sup>C.
44. (New) (Currently amended) A polythiophene which comprise recurring units of the formula (IV),





(IV)


wherein

A is a C<sub>1</sub>-C<sub>5</sub>-alkylene radical which is substituted at any point by a linker L and optionally bears further substituents,


L is a methylene group,

p is 0 or an integer from 1 to 6,

M is an n-functional group of the formula (II-a) or (II-b),
















(II-a)




(II-b)

wherein


X<sup>1</sup>, X<sup>2</sup> and X<sup>3</sup> are substituted or unsubstituted structures selected independently from the group consisting of





and

$Z^1$  and  $Z^2$  are structures selected independently from the group consisting of



wherein

$R^x$  and  $R^y$  are each, independently of one another, H, substituted or unsubstituted  $C_1$ - $C_{22}$ -alkyl,  $C_1$ - $C_{22}$ -haloalkyl,  $C_1$ - $C_{22}$ -alkenyl,  $C_1$ - $C_{22}$ -alkoxy,  $C_1$ - $C_{22}$ -thioalkyl,  $C_1$ - $C_{22}$ -iminoalkyl,  $C_1$ - $C_{22}$ -alkoxycarbonyl,  $C_1$ - $C_{22}$ -alkoxycarbonyloxy, a radical of an aliphatic  $C_1$ - $C_{22}$ -alkanecarboxylic acid or of acrylic acid, halogen, pseudohalogen,  $NO_2$ , a carboxyl group or a hydroxy group,

$h$  is an integer from 1 to 10,

w is an integer from 1 to 5,

x, y and z are each, independently of one another, 0 or 1, and


n is 1 or 2, where when n is 1, the group of the formula (II-a) or (II-b) bears a terminal group F at the linkage points denoted by \*,

wherein

F is substituted or unsubstituted C<sub>1</sub>-C<sub>22</sub>-alkyl, C<sub>1</sub>-C<sub>22</sub>-haloalkyl, C<sub>1</sub>-C<sub>22</sub>-alkenyl, C<sub>1</sub>-C<sub>22</sub>-alkoxy, C<sub>1</sub>-C<sub>22</sub>-thioalkyl, C<sub>1</sub>-C<sub>22</sub>-iminoalkyl, C<sub>1</sub>-C<sub>22</sub>-alkoxycarbonyl, C<sub>1</sub>-C<sub>22</sub>-alkoxycarbonyloxy, a radical of an aliphatic C<sub>1</sub>-C<sub>22</sub>-alkanecarboxylic acid or of acrylic acid, halogen, pseudohalogen, a nitro (NO<sub>2</sub>) group, a carboxyl group, a sulphonic acid group or sulphonate group or a hydroxy group,

n is an integer from 1 to 8 and

B is a bridging group of the formula (B)



wherein

q is 0 or 1,

r and s are each 0 or 1, with the proviso that when r is 1, s is 0 and vice versa or both are optionally 0,


t is 0 or 1,

Sp is a spacer selected from the group consisting of substituted and unsubstituted linear or cyclic C<sub>1</sub>-C<sub>20</sub>-alkylene groups, C<sub>5</sub>-C<sub>20</sub>-arylene groups, C<sub>2</sub>-C<sub>20</sub>-heteroarylene groups in which from one to three heteroatoms selected from the group consisting of N, O and S can additionally be present in the heteroaromatic ring or ring system, C<sub>6</sub>-C<sub>20</sub>-aralkylene groups, C<sub>2</sub>-C<sub>200</sub>-oligoether and -polyether groups,


m is 0 or 1,

Q is O, S or NH,

with the proviso that said polythiophenes do not contain recurring units of the formula (ii)




45. (New) Process for preparing polythiophenes, comprising oxidatively polymerizing electrochemically compounds of the formula (I),



where

- A is a  $C_1$ - $C_5$ -alkylene radical which is substituted at any point by a linker L and optionally bears further substituents,
- L is a methylene group,
- p is 0 or an integer from 1 to 6,
- n is an integer from 1 to 8 and
- B is a bridging group of the formula (B)



wherein

q is 0 or 1,

r, s are each 0 or 1, with the proviso that when r is 1, s is 0 and vice versa or both are optionally 0,


t is 0 or 1,

Sp is a spacer selected from the group consisting of substituted and unsubstituted linear or cyclic C<sub>1</sub>-C<sub>20</sub>-alkylene groups, C<sub>5</sub>-C<sub>20</sub>-arylene groups, C<sub>2</sub>-C<sub>20</sub>-hetero-arylene groups in which from one to three heteroatoms selected from the group consisting of N, O and S can additionally be present in the heteroaromatic ring or ring system, C<sub>6</sub>-C<sub>20</sub>-aralkylene groups, C<sub>2</sub>-C<sub>200</sub>-oligoether and -polyether groups,


m is 0 or 1,

Q is O, S or NH, and

M is an n-functional group of the formula (II-a) or (II-b),















(II-a)




(II-b)

wherein

X<sup>1</sup>, X<sup>2</sup> and X<sup>3</sup> are substituted or unsubstituted structures selected independently from the group consisting of








and

$Z^1$  and  $Z^2$  are structures selected independently from the group consisting of



wherein

$\text{R}^x$  and  $\text{R}^y$  are each, independently of one another, H, substituted or unsubstituted  $\text{C}_1$ - $\text{C}_{22}$ -alkyl,  $\text{C}_1$ - $\text{C}_{22}$ -haloalkyl,  $\text{C}_1$ - $\text{C}_{22}$ -alkenyl,  $\text{C}_1$ - $\text{C}_{22}$ -alkoxy,  $\text{C}_1$ - $\text{C}_{22}$ -thioalkyl,  $\text{C}_1$ - $\text{C}_{22}$ -iminoalkyl,  $\text{C}_1$ - $\text{C}_{22}$ -alkoxycarbonyl,  $\text{C}_1$ - $\text{C}_{22}$ -alkoxycarbonyloxy, a radical of an aliphatic  $\text{C}_1$ - $\text{C}_{22}$ -alkanecarboxylic acid or of acrylic acid, halogen, pseudohalogen,  $\text{NO}_2$ , a carboxyl group or a hydroxy group,

$h$  is an integer from 1 to 10,

n is 1 or 2, where when n is 1, the group of the formula (II-a) or (II-b) bears a terminal group F at the linkage points denoted by \*,

F is substituted or unsubstituted C<sub>1</sub>-C<sub>22</sub>-alkyl, C<sub>1</sub>-C<sub>22</sub>-haloalkyl, C<sub>1</sub>-C<sub>22</sub>-alkenyl, C<sub>1</sub>-C<sub>22</sub>-alkoxy, C<sub>1</sub>-C<sub>22</sub>-thioalkyl, C<sub>1</sub>-C<sub>22</sub>-iminoalkyl, C<sub>1</sub>-C<sub>22</sub>-alkoxycarbonyl, C<sub>1</sub>-C<sub>22</sub>-alkoxycarbonyloxy, a radical of an aliphatic C<sub>1</sub>-C<sub>22</sub>-alkanecarboxylic acid or of acrylic acid, halogen, pseudohalogen, a nitro (NO<sub>2</sub>) group, a carboxyl group, a sulphonic acid group or sulphonate group or a hydroxy group

N#Cc1ccc(cc1)-c2ccc(cc2)Oc3ccc(cc3)OCCOC4C5C(C=C4)OC6C(=C5)SC=C6